

```
In [ ]: cd matlab
        pwd
```

Fourier Transforms for Circuit and LTI Systems Analysis

Scope and Background Reading

This session we will apply what we have learned about Fourier transforms to some typical circuit problems. After a short introduction, this session will be an examples class.

The material in this presentation and notes is based on Chapter 8 (Starting at Section 8.8) of [Steven T. Karris, Signals and Systems: with Matlab Computation and Simulink Modelling, 5th Edition.](#) (<https://ebookcentral.proquest.com/lib/swansea-ebooks/reader.action?ppg=304&docID=3384197&tm=1518713030874>) from the **Required Reading List**. I also used Chapter 5 of [Benoit Boulet, Fundamentals of Signals and Systems](#) (<https://ebookcentral.proquest.com/lib/swansea-ebooks/reader.action?ppg=212&docID=3135971&tm=1518713118808>) from the **Recommended Reading List**.

Agenda

- The system function
- Examples

The System Function

System response from system impulse response

Recall that the convolution integral of a system with impulse response $h(t)$ and input $u(t)$ is

$$h(t) * u(t) = \int_{-\infty}^{\infty} h(t - \tau)u(\tau) d\tau.$$

We let

$$g(t) = h(t) * u(t)$$

Then by the time convolution property

$$h(t) * u(t) = g(t) \Leftrightarrow G(\omega) = H(\omega) \cdot U(\omega)$$

The System Function

We call $H(\omega)$ the *system function*.

We note that the system function $H(\omega)$ and the impulse response $h(t)$ form the Fourier transform pair

$$h(t) \Leftrightarrow H(\omega)$$

Obtaining system response


If we know the impulse response $h(t)$, we can compute the system response $g(t)$ of any input $u(t)$ by multiplying the Fourier transforms of $H(\omega)$ and $U(\omega)$ to obtain $G(\omega)$. Then we take the inverse Fourier transform of $G(\omega)$ to obtain the response $g(t)$.

1. Transform $h(t) \rightarrow H(\omega)$
2. Transform $u(t) \rightarrow U(\omega)$
3. Compute $G(\omega) = H(\omega) \cdot U(\omega)$
4. Find $\mathcal{F}^{-1} \{G(\omega)\} \rightarrow g(t)$


Examples

Example 1

Karris example 8.8: for the linear network shown below, the impulse response is $h(t) = 3e^{-2t}$. Use the Fourier transform to compute the response $y(t)$ when the input $u(t) = 2[u_0(t) - u_0(t - 3)]$. Verify the result with Matlab.



Solution



Matlab verification

```
In [2]: syms t w
        U1 = fourier(2*heaviside(t),t,w)
```

```
U1 =
2*pi*dirac(w) - 2i/w
```

```
In [3]: H = fourier(3*exp(-2*t)*heaviside(t),t,w)
```

```
H =
3/(2 + w*1i)
```

```
In [4]: Y1=simplify(H*U1)
```

```
Y1 =
3*pi*dirac(w) - 6i/(w*(2 + w*1i))
```

```
In [5]: y1 = simplify(ifourier(Y1,w,t))
```

```
y1 =
(3*exp(-2*t)*(sign(t) + 1)*(exp(2*t) - 1))/2
```

Get y2

Substitute $t - 3$ into t .

```
In [6]: y2 = subs(y1,t,t-3)
```

```
y2 =
(3*exp(6 - 2*t)*(sign(t - 3) + 1)*(exp(2*t - 6) - 1))/2
```


```
In [7]: y = y1 - y2
```

y =

$$(3 \cdot \exp(-2 \cdot t) \cdot (\text{sign}(t) + 1) \cdot (\exp(2 \cdot t) - 1)) / 2 - (3 \cdot \exp(6 - 2 \cdot t) \cdot (\text{sign}(t - 3) + 1) \cdot (\exp(2 \cdot t - 6) - 1)) / 2$$

Plot result

```
In [8]: ezplot(y)
title('Solution to Example 1')
ylabel('y(y)')
xlabel('t [s]')
grid
```



See [ft3_ex1.m \(matlab/ft3_ex1.m\)](http://localhost:8890/nbconvert/html/week7/ft3.ipynb?download=false)

Result is equivalent to:


$$y = 3 \cdot \text{heaviside}(t) - 3 \cdot \text{heaviside}(t - 3) + 3 \cdot \text{heaviside}(t - 3) \cdot \exp(6 - 2 \cdot t) - 3 \cdot \exp(-2 \cdot t) \cdot \text{heaviside}(t)$$

Which after gathering terms gives

$$y(t) = 3(1 - 3e^{-2t})u_0(t) - 3(1 - 3e^{-2(t-3)})u_0(t - 3)$$

Example 2

Karris example 8.9: for the circuit shown below, use the Fourier transform method, and the system function $H(\omega)$ to compute $V_L(t)$. Assume $i_L(0^-) = 0$. Verify the result with Matlab.



Solution



Matlab verification

```
In [9]: syms t w  
H = j*w/(j*w + 2)
```

H =

$$(w*1i)/(2 + w*1i)$$

```
In [10]: Vin = fourier(5*exp(-3*t)*heaviside(t),t,w)
```

Vin =

$$5/(3 + w*1i)$$

```
In [11]: Vout=simplify(H*Vin)
```

Vout =

$$(w*5i)/((2 + w*1i)*(3 + w*1i))$$


```
In [12]: vout = simplify(iffourier(Vout,w,t))
```

vout =

$$-(5*\exp(-3*t))*(\text{sign}(t) + 1)*(2*\exp(t) - 3))/2$$

Plot result


```
In [13]: ezplot(vout)
         title('Solution to Example 2')
         ylabel('v_{out}(t) [V]')
         xlabel('t [s]')
         grid
```



See [ft3_ex2.m \(matlab/ft3_ex2.m\)](#)

Result is equivalent to:

$$v_{out} = -5 \cdot \exp(-3 \cdot t) \cdot \text{heaviside}(t) \cdot (2 \cdot \exp(t) - 3)$$

Which after gathering terms gives



$$v_{out} = 5 \left(3e^{-3t} - 2e^{-2t} \right) u_0(t)$$

Example 3


Karris example 8.10: for the linear network shown below, the input-output relationship is:

$$\frac{d}{dt}v_{\text{out}} + 4v_{\text{out}} = 10v_{\text{in}}$$

where $v_{\text{in}} = 3e^{-2t}$. Use the Fourier transform method, and the system function $H(\omega)$ to compute the output v_{out} . Verify the result with Matlab.



Solution



Matlab verification

```
In [14]: syms t w  
H = 10/(j*w + 4)
```

H =

$10/(4 + w*1i)$

```
In [15]: Vin = fourier(3*exp(-2*t)*heaviside(t),t,w)
```

Vin =

$3/(2 + w*1i)$

```
In [16]: Vout=simplify(H*Vin)
```

Vout =

$$30/((2 + w*1i)*(4 + w*1i))$$


```
In [17]: vout = simplify(ifourier(Vout,w,t))
```

vout =

$$(15*\exp(-4*t)*(sign(t) + 1)*(exp(2*t) - 1))/2$$

Plot result

```
In [18]: ezplot(vout)
title('Solution to Example 2')
ylabel('v_{out}(t) [V]')
xlabel('t [s]')
grid
```



See [ft3_ex3.m \(matlab/ft3_ex3.m\)](#)

Result is equivalent to:

$$15 * \exp(-4 * t) * \text{heaviside}(t) * (\exp(2 * t) - 1)$$

Which after gathering terms gives


$$v_{\text{out}}(t) = 15 (e^{-2t} - e^{-4t}) u_0(t)$$

Example 4

Karris example 8.11: the voltage across a 1Ω resistor is known to be $V_R(t) = 3e^{-2t}u_0(t)$. Compute the energy dissipated in the resistor for $0 < t < \infty$, and verify the result using Parseval's theorem. Verify the result with Matlab.

Note from [tables of integrals \(http://en.wikipedia.org/wiki/Lists_of_integrals\)](http://en.wikipedia.org/wiki/Lists_of_integrals)

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \arctan \frac{x}{a} + C.$$



Solution



Matlab verification

In [19]: `syms t w`

Calculate energy from time function

```
In [20]: Vr = 3*exp(-2*t)*heaviside(t);
R = 1;
Pr = Vr^2/R
Wr = int(Pr,t,0,inf)
```

Pr =

$9 \cdot \exp(-4 \cdot t) \cdot \text{heaviside}(t)^2$

Wr =

$9/4$

Calculate using Parseval's theorem

```
In [21]: Fw = fourier(Vr,t,w)
```

Fw =

$3/(2 + w \cdot 1i)$

```
In [22]: Fw2 = simplify(abs(Fw)^2)
```

Fw2 =

$9/\text{abs}(2 + w \cdot 1i)^2$

```
In [23]: Wr=2/(2*pi)*int(Fw2,w,0,inf)
```

Wr =

$(51607450253003931 \cdot \pi) / 72057594037927936$

See [ft3_ex4.m \(matlab/ft3_ex4.m\)](#)